

# Examples and Figures from Wang et al. Journal of Circadian Rhythms 2011

Carlos Gonzalez, Berkley Shands, Elena Deych, William Shannon

January 15, 2016

First we need to load the Actigraphy package:

```
> library(Actigraphy)
> library(lattice)
```

## 1 Figure 1(a)

```
> ### Load Data
> data(weekday)
> ### Data Management
> data2 <- NULL
> data2$act <- weekday[,3]
> data2$t <- weekday[,2]
> data2$day <- weekday[,1]
> data2$date <- factor(data2$day, levels=c("Monday", "Tuesday", "Wednesday", "Thur
> ### Plot Options and Parameters
> lb <- c("Midnight", "6AM", "Noon", "6PM", "Midnight")
> L <- 1440
> xat <- c(0, L/4, L/2, 3*L/4, L)
> ### Plot Figure
> xyplot(act ~ t | date, data=data2, as.table=TRUE,
+         main="Subject 002 Activity from Monday to Friday",
+         scales=list(x=list(at=xat, labels=lb)), cex.main=0.5,
+         layout=c(1, 5, 1), xlim=c(0, L), xlab="(a)",
+         ylab="Activity", panel=function(x,y) {
+           fbase <- create.fourier.basis(rangeval=c(0,L), nbasis=9)
+           fpar <- fdPar(fbase)
+           fd <- smooth.basis(c(1:L), y, fpar)
+           panel.xyplot(x, y, type="h")
+         })
```

## 2 Figure 2

```
> ### Load Data
> data(act_8pt)
> data(clinic_8pt)
```

```

>     ### Plot Options and Parameters
>     lb <- c("Midnight", "6AM", "Noon", "6PM", "Midnight")
>     L <- 1440
>     xat <- c(0, L/4, L/2, 3*L/4, L)
>     matchid <- fda.matchid(act_8pt[,-1], clinic_8pt, type="factor", grouplab=c("AHI"
>     idhigh <- paste("Subj", colnames(matchid$mat)[matchid$cov$"NO AHI" != 1])
>     idlow <- paste("Subj", colnames(matchid$mat)[matchid$cov$"NO AHI" == 1])
>     idorder <- c(idhigh, idlow)
>     datavec <- matchid$mat
>     dim(datavec) <- NULL
>     datanew <- data.frame(y=datavec, id=rep(paste("Subj", colnames(matchid$mat)), ea
>     datanew$id <- factor(datanew$id, idorder)
>     ### Plot Figure
>     xyplot(y~t|id, data=datanew, as.table=TRUE,
+           main="Circadian Activity from 8 Subjects",
+           ylab="Activity", xlab="", cex.main=.7,
+           scales=list(x=list(at=xat, labels=lb)), cex=.05,
+           type="p", layout=c(4, 2, 1), ylim=c(0, 1200),
+           xlim=c(0, L), panel=function(x,y) {
+               fbase <- create.fourier.basis(rangeval=c(0, L), nbasis=9)
+               fpar <- fdPar(fbase)
+               sm <- smooth.basis(c(1:L), y, fpar)
+               panel.xyplot(x, y, col=1, cex=0.1)
+               panel.lines(predict(sm$fd,c(1:L)), col=2, lwd=3)
+           })

```

### 3 Figure 3

```

>     ### Load Data
>     data(act_8pt)
>     data(clinic_8pt)
>     ahidatav2 <- fda.matchid(act_8pt[,-1], clinic_8pt, type="factor", grouplab=c("AHI"
>     tempv2 <- ahidatav2[[2]]
>     tempv2[,3] <- ifelse(tempv2[,3] == 0, -1, 1)
>     ahidatav2$cov <- data.frame(id=tempv2$id, mean=1, ahi=tempv2[,3])
>     colv2 <- ifelse(tempv2[,3] == -1, 4, 2)
>     smoothDatav2 <- fda.smoothdata(ahidatav2)
>     geftahiv2 <- flm_cate(smoothDatav2)
>     meanefv2 <- geftahiv2$freg$betaestlist[[1]]
>     ahiefv2 <- geftahiv2$freg$betaestlist[[2]]
>     ### Plot Options and Parameters
>     L <- 1440
>     xat <- c(0, L/4, L/2, 3*L/4, L)
>     lb <- c("Midnight", "6AM", "Noon", "6PM", "Midnight")
>     ### Plot Figure
>     par(mfrow=c(2,1), mar=c(4,4,3,1))
>     plot(0, 0, xlim=c(0,L), ylim=c(0,1200), xaxt="n", xlab="(a)", ylab="Acitivity",
>     for(i in 1:8)
+         lines(predict(smoothDatav2$fd$fd, c(1:L))[,i], col=colv2[i])

```

```

>     ### Plot the group mean activities
>     lines(meanefv2$fd-ahiefv2$fd, col=4, lwd=3)
>     lines(meanefv2$fd+ahiefv2$fd, col=2, lwd=3)
>     ### Plot the overall mean
>     lines(meanefv2$fd, col=1, lwd=3)
>     ### Add the axis and legend to finish the plot
>     axis(1, at=xat, labels=lb)
>     legend("topleft", c("AHI High Curves", "AHI High Mean", "AHI Low Curves", "AHI L
+         lty=1, col=c(4,4,2,2,1), lwd=c(1,3,1,3,3), cex=.8)
>     ### F Test
>     cov2 <- smoothDatav2$cov[, -1]
>     grp2 <- ncol(cov2)
>     fd <- smoothDatav2$fd
>     L <- length(fd$argvals)
>     npt <- ncol(fd$y)
>     fbase <- create.fourier.basis(rangeval=c(0, 1440), nbasis=9)
>     fpar <- fdPar(fbase)
>     xfdlist <- vector("list", grp2)
>     xfdlist[[1]] <- cov2[, 1] + 0
>     for(i in 2:grp2)
+         xfdlist[[i]] <- cov2[, i] + 0
>     betalist <- xfdlist
>     for(i in 1:grp2)
+         betalist[[i]] <- fpar
>     freg2 <- fRegress(fd$fd, xfdlist, betalist)
>     preact2 <- predict(freg2$yhatfdobj, c(1:L))
>     resid2 <- fd$y - preact2[, 1:npt]
>     sigma2 <- cov(t(resid2))
>     fregstd2 <- fRegress.stderr(freg2, fd$y2cMap, sigma2)
>     Fratio <- Ftest(fd$fd, xfdlist, betalist, argvals = c(1:1440), nperm=10, xaxt="
>     axis(1, at=xat, labels=lb)

```