

Package ‘perccalc’

September 14, 2017

Title Estimate Percentiles from an Ordered Categorical Variable

Version 1.0.0

Description An implementation of two functions that estimate values for percentiles from an ordered categorical variable as described by Reardon (2011, isbn:978-0-87154-372-1). One function estimates percentile differences from two percentiles while the other returns the values for every percentile from 1 to 100.

Depends R (>= 3.4.0)

License MIT + file LICENSE

Encoding UTF-8

LazyData true

RoxygenNote 6.0.1

Imports magrittr, dplyr, purrr, stats, tibble, broom, tidyr, multcomp, haven, tidyverse, ggplot2, devtools, MASS

Suggests spelling, knitr, rmarkdown

VignetteBuilder knitr

NeedsCompilation no

Author Jorge Cimentada [aut, cre]

Maintainer Jorge Cimentada <cimentadaj@gmail.com>

Repository CRAN

Date/Publication 2017-09-14 17:47:40 UTC

R topics documented:

perc_diff	2
perc_dist	3
Index	5

perc_diff	<i>Calculate percentile differences from an ordered categorical variable and a continuous variable.</i>
-----------	---

Description

Calculate percentile differences from an ordered categorical variable and a continuous variable.

Usage

```
perc_diff(data_model, categorical_var, continuous_var, weights = NULL,  
          percentiles = c(90, 10))
```

Arguments

data_model	A data frame with at least the categorical and continuous variables from which to estimate the percentile differences
categorical_var	The bare unquoted name of the categorical variable. This variable SHOULD be an ordered factor. If not, the function will stop.
continuous_var	The bare unquoted name of the continuous variable from which to estimate the percentiles
weights	The bare unquoted name of the optional weight variable. If not specified, then estimation is done without weights
percentiles	A numeric vector of two numbers specifying which percentiles to subtract

Value

A vector with the percentile difference and its associated standard error

Examples

```
library(dplyr)  
  
set.seed(23131)  
N <- 1000  
K <- 20  
  
toy_data <- tibble::tibble(id = 1:N,  
  score = rnorm(N, sd = 2),  
  type = rep(paste0("inc", 1:20), each = N/K),  
  wt = 1)  
  
# perc_diff(toy_data, type, score)  
# type is not an ordered factor!  
  
toy_data <-
```

```
toy_data %>%
mutate(type = factor(type, levels = unique(type), ordered = TRUE))

perc_diff(toy_data, type, score, percentiles = c(90, 10))
perc_diff(toy_data, type, score, percentiles = c(50, 10))

perc_diff(toy_data, type, score, weights = wt, percentiles = c(30, 10))
```

perc_dist	<i>Calculate a distribution of percentiles from an ordered categorical variable and a continuous variable.</i>
-----------	--

Description

Calculate a distribution of percentiles from an ordered categorical variable and a continuous variable.

Usage

```
perc_dist(data_model, categorical_var, continuous_var, weights = NULL)
```

Arguments

data_model	A data frame with at least the categorical and continuous variables from which to estimate the percentiles
categorical_var	The bare unquoted name of the categorical variable. This variable SHOULD be an ordered factor. If not, the function will stop.
continuous_var	The bare unquoted name of the continuous variable from which to estimate the percentiles
weights	The bare unquoted name of the optional weight variable. If not specified, then estimation is done without weights

Value

A data frame with the scores and standard errors for each percentile

Examples

```
library(dplyr)

set.seed(23131)
N <- 1000
K <- 20

toy_data <- tibble::tibble(id = 1:N,
```

```
score = rnorm(N, sd = 2),
type = rep(paste0("inc", 1:20), each = N/K,
wt = 1)

# perc_dist(toy_data, type, score)
# type is not an ordered factor!

toy_data <-
  toy_data %>%
  mutate(type = factor(type, levels = unique(type), ordered = TRUE))

perc_dist(toy_data, type, score)
```

Index

perc_diff, 2
perc_dist, 3