

# Package ‘CATTexact’

September 15, 2017

**Type** Package

**Title** Computation of the p-Value for the Exact Conditional  
Cochran-Armitage Trend Test

**Version** 0.1.0

## Description

Provides functions for computing the one-sided p-values of the Cochran-Armitage trend test statistic for the asymptotic and the exact conditional test. The computation of the p-value for the exact test is performed using an algorithm following an idea by Mehta, et al. (1992) <doi:10.2307/1390598>.

**Depends** R (>= 3.4.1)

**License** GPL-2 | GPL-3

**LazyData** TRUE

**RoxygenNote** 6.0.1

**Suggests** testthat

**NeedsCompilation** no

**Author** Dominic Edelmann [aut, cre]

**Maintainer** Dominic Edelmann <dominic.edelmann@dkfz-heidelberg.de>

**Repository** CRAN

**Date/Publication** 2017-09-15 11:34:22 UTC

## R topics documented:

catt_asy . . . . .	2
catt_exact . . . . .	3

<b>Index</b>	<b>5</b>
--------------	----------

---

`catt_asy`*Asymptotic Cochran-Armitage trend test*

---

**Description**

`catt_asy` calculates the Cochran-Armitage trend test statistic (Cochran (1954), Armitage (1955)) and the one-sided p-value for the corresponding asymptotic test. The exact form of used test statistic can be found in the paper by Portier and Hoel (1984).

**Usage**

```
catt_asy(dose.ratings, totals, cases)
```

**Arguments**

<code>dose.ratings</code>	A vector of dose ratings, the i-th entry corresponds to the dose-rating of the i-th group. This vector must be strictly monotonically increasing
<code>totals</code>	The vector of total individuals per group, the i-th entry corresponds to the total number of individuals in the i-th group
<code>cases</code>	The vector of incidences per groups, the i-th entry corresponds to the number of incidences in the i-th group

**Value**

A list containing the value of the Cochran-Armitage Trend Test Statistic and its asymptotic p-value.

**References**

Armitage, P. Tests for linear trends in proportions and frequencies. *Biometrics*, 11 (1955): 375-386.

Cochran, W. G. Some methods for strengthening the common  $\chi^2$  tests, *Biometrics*. 10 (1954): 417-451.

Portier, C., and Hoel D. Type 1 error of trend tests in proportions and the design of cancer screens. *Communications in Statistics-Theory and Methods*, 13 (1984): 1-14.

**Examples**

```
d <- c(1,2,3,4)
n <- rep(20,4)
r <- c(1,4,3,8)

catt_asy(d, n, r)
```

---

`catt_exact`*Conditional exact Cochran-Armitage trend test*

---

**Description**

`catt_exact` calculates the Cochran-Armitage trend test statistic (Cochran (1954), Armitage (1955)) and the one-sided p-value for the corresponding conditional exact test. The conditional exact test has been established by Williams (1988). The computation of its p-value is performed using an algorithm following an idea by Mehta, et al. (1992).

**Usage**

```
catt_exact(dose.ratings, totals, cases)
```

**Arguments**

<code>dose.ratings</code>	A vector of dose ratings, the i-th entry corresponds to the dose-rating of the i-th group. This vector must be strictly monotonically increasing
<code>totals</code>	The vector of total individuals per group, the i-th entry corresponds to the total number of individuals in the i-th group.
<code>cases</code>	The vector of incidences per groups, the i-th entry corresponds to the number of incidences in the i-th group.

**Value**

A list containing the value of the Cochran-Armitage Trend Test Statistic, its exact and asymptotic p-value.

**References**

Armitage, P. Tests for linear trends in proportions and frequencies. *Biometrics*, 11 (1955): 375-386.

Cochran, W. G. Some methods for strengthening the common  $\chi^2$  tests, *Biometrics*. 10 (1954): 417-451.

Mehta, C. R., Nitin P., and Pralay S. Exact stratified linear rank tests for ordered categorical and binary data. *Journal of Computational and Graphical Statistics*, 1 (1992): 21-40.

Portier, C., and Hoel D. Type 1 error of trend tests in proportions and the design of cancer screens. *Communications in Statistics-Theory and Methods*, 13 (1984): 1-14.

Williams, D. A. Tests for differences between several small proportions. *Applied Statistics*, 37 (1988): 421-434.

**Examples**

```
d <- c(1,2,3,4)
n <- rep(20,4)
r <- c(1,4,3,8)

catt_exact(d, n, r)
```

# Index

catt\_asy, 2  
catt\_exact, 3